NEW HOST RECORDS IN THE BRUCHIDAE (COLEOPTERA)

CLARENCE DAN JOHNSON

Department of Biological Sciences, Northern Arizona University, Flagstaff, AZ 86011

ABSTRACT

Eight species of the genus Amblycerus are reported to feed in seeds of species of the genera Caesalpinia, Cassia, Apeiba, and Combretum. Meibomeus musculus (Say) feeds in species of Desmodium and Lespedeza; seven species of Merobruchus feed in seeds of species of Albizia, Lysiloma, Pithecellobium, Acacia, and Leucaena; Mimosestes nubigens (Motschulsky) feeds in seeds of Acacia; Sennius morosus (Sharp) feeds in seeds of Cassia durangensis Rose; six species of Stator feed in seeds of species of Acacia, Calliandra, Lysiloma, Leucaena, and Mimosa; and four species of Zabrotes feed in seeds of Cassia and Vicia. All of the plants are in the family Leguminosae except for Apeiba (Tiliaceae) and Combretum (Combretaceae).

To make them available for current review taxonomic and ecological studies currently in progress, new host records for the genera Amblycerus, Meibomeus, Merobruchus, Mimosestes, Sennius, Stator, and Zabrotes are presented below. The species designated by number are either new or too variable to be assigned a name prior to a thorough revision of those genera. J. M. Kingsolver is currently revising Amblycerus and Merobruchus and will assign names to these numbers when the revisions are published. Unless otherwise indicated, all plants are in the family Leguminosae.

ACKNOWLEDGMENTS

I am grateful to J. M. Kingsolver for verifications of identifications of bruchids; D. E. Bright and E. Becker for making available the host records of L. J. Bottimer; R. Liesner, T. Croat, J. Strother, and R. Barneby for plant determinations; and the Johnsons, Debbie, Jack, Margaret, and Rod, for help in collecting seeds and plants.

The Insect Identification and Beneficial Insect Introduction Institute, U.S. Department of Agriculture, provided funds for collecting host data under Grants 12-14-100-9187(33) and 12-14-100-9970(33). National Science Foundation Grant DEB78-05962 provided funds for compilation and analysis of data.

In the text of this paper C. D. Johnson is abbreviated CDJ and L. J. Bottimer LJB.

Amblycerus sp. #1

1. Caesalpinia sclerocarpa Standley: Mexico. Sinaloa: near beach, 6 mi N Mazatlan, 26-II-73 (CDJ #215-73).

Amblycerus sp. #2

1. Cassia berlandieri Bentham: Mexico. Nayarit: 29 mi SW Compostela, 1-III-73 (CDJ #302-73).

Amblycerus sp. #3

1. Cassia bicapsularis Linnaeus: Mexico. Sinaloa: 3 mi NE Los Mochis, 20-VIII-65 (CDJ). Oaxaca: 5 mi E Temascal, 24-VI-64 (D. H. Janzen).

2. C. occidentalis Linnaeus: Mexico. Navarit: 4 mi E San Blas, 27-VIII-65 (CDJ).

Amblycerus sp. #4

1. Cassia emarginata Linnaeus: Mexico. Colima: 11 mi NE Armeria, 7-III-73 (CDJ #390-73).

Amblycerus sp. #5

1. Cassia hintoni Sandwith: Mexico. Jalisco: 22 mi N Barra de Navidad, 2-I-73 (CDJ #28-73).

Amblycerus sp. #6

1. Cassia multijuga Rich.: Florida. Dade Co.: Homestead, 8-IV-60 (LJB #106f).

Amblycerus sp. #7

1. Apeiba membranacea Spruce ex Benth.: Panama (LJB). (Tiliaceae)

Amblycerus sp. #8

1. Combretum fruticosum (Loefl.) Stuntz: Colima: 7 mi E Manzanillo, 7-III-73 (CDJ #408-73). Jalisco: 10 mi S Puerto Vallarta, 9-III-73 (CDJ #460-73). (Combretaceae)

Meibomeus musculus (Say)

1. Desmodium canescens DC: Florida. Polk Co.: Lake Alfred, 6-VII-28 (LJB #55a8).

2. D. purpureum Hook. & Arn.: Florida. Alachua Co.: Gainesville, 4-VIII-37 (A. N. Tissot; LJB #761).

3. D. tenuifolium Torr. & Gray: Florida. Polk Co.: Dundee, VII-28 (LJB #55a5).

4. D. triflorum DC: Florida. Orange Co.: 9 mi S Orlando, 25-VI-28 (LJB #55b2). 5. Lespedeza hirta Ell.: New Jersey. Camden Co.: Haddon Heights, 30-VII-36 (LJB #741).

Merobruchus insolitus (Sharp)

1. Albizia lebbek (L.) Bentham: Mexico. Baja California Sur: San Jose Del Cabo, 27-XII-75 (CDJ #129-75).

2. A. occidentalis Brandegee: Mexico. Baja California Sur: 700', 12 mi N San Jose

Del Cabo, 28-XII-75 (CDJ #141-75).

3. A. sinaloensis Britt. & Rose: Mexico. Sonora: Alamos, 30 December 1975

(CDJ # 146-75).

4. Lysiloma divaricata (Jacq.) Macbride: Mexico. Sinaloa: 30 mi S Guamuchil, 7-I-73 (CDJ #137-73). Jalisco: ca 4000′, 3 mi S Tecolatlan, 1-I-73 (CDJ #3-73). Sonora: 10 mi N Guaymas, 3-I-78 (CDJ #220-78).

5. Lysiloma thornberi Britt. & Rose: Arizona. Pima Co.: Chimenea Cyn., Rincon

Mts., 14-X-76 (CDJ # #37-76 & 38-76).

6. Pithecellobium aff. mangense (Jacq.) Macbride: Mexico. Nayarit: 2 mi NW Acaponeta, 10-III-73 (CDJ #482-73).

7. P. sonorae S. Wats.: Mexico. Sonora: 9 mi E Navojoa, 30-XII-75 (CDJ # 151-75).

M. julianus (Horn)

1. Acacia roemeriana Scheele: Texas. Menard Co.: Menard, 27-VI-46 (LJB # 86h & 86i). Kimble Co.: Junction, 17-VI-56 (LJB # 97i).

M. knulli (White)

1. Lysiloma thornberi Britt. & Rose: Arizona. Pima Co.: Chimenea Cyn., Rincon Mts., 14-X-76 (CDJ # #37-76 & 38-76).

M. placidus (Horn)

1. Acacia texensis Torr. & Gray: Texas. Big Bend National Park: Oak Spring, 1-V-59 (LJB #104q).

Merobruchus sp. #1

1. Acacia coulteri Bentham: Mexico. Sonora: 8 mi E Navojoa, 24-II-73 (CDJ #179-73); Lake Mocuzari, 29-XII-77 (CDJ #191-77); 9 mi E Navojoa, 30-XII-75 (CDJ #153-75).

Merobruchus sp. #2

1. Pithecellobium aff. mangense (Jacq.) Macbride: Mexico. Sinaloa: 38 mi S Culiacan, 25-II-73 (CDJ #196-73). Nayarit: 2 mi NW Acaponeta, 10-III-73 (CDJ #482-73).

2. P. undulatum (Britt. & Rose) Gentry: Mexico. Sonora: 13 mi NW Alamos & 4 mi NW Alamos, 23-XII-76 (CDJ ##165-76 & 175-76); 5 mi W Alamos, 26-XII-76 (CDJ #186-76); 6 mi W Alamos, 27-XII-77 (CDJ #168-77); 2 mi E Alamos, 29-XII-77 (CDJ #189-77).

Merobruchus sp. #3

- 1. Acacia aff. riparioides (Britt. & Rose) Standley: Mexico. Michoacan: 5300′, 16 mi E Jiquilpan, 9-VII-68 (CDJ # 256-68).
- 2. Leucaena guatemalensis Britt. & Rose: Mexico. Colima: 3 mi S Colima, 7-III-73 (CDJ #373-73).
- 3. Lysiloma desmostachys Bentham: Mexico. Colima: 7 mi S Colima, 7-III-73 (CDJ # 380-73).

Merobruchus sp. #4

1. Acacia sp.: Mexico. Jalisco: 56 mi S Puerto Vallarta, 9-III-73 (CDJ #446-73).

Mimosestes nubigens (Motschulsky)

1. Acacia tortuosa Willd.: Mexico. Sonora: 13 mi NW Navojoa, 30-XII-77 (CDJ #206-77).

Sennius medialis (Sharp)

1. Cassia occidentalis Linnaeus: Mexico. Michoacan: Patzcuaro, 9-VIII-49 (LJB #125f)

Sennius morosus (Sharp)

1. Cassia durangensis Rose: Texas: Santa Helena, 1900', 10-II-26 (LJB #54rl). Big Bend National Park: Boquillas Canyon, 5-V-56 (LJB #96v); Hot Springs, 19-VI-57 (LJB #100q).

Stator chihuahua Johnson and Kingsolver

- 1. Acacia angustissima (Mill.) Kuntze: Mexico. Sonora: ca 2800', 8 mi S Benjamin Hill, 28-XII-76 (CDJ #221-76); ca 2400', 15 mi S Benjamin Hill, 3-I-78 (CDJ #233-78).
- 2. A. constricta Bentham: Arizona. Pima Co.: 1 mi S Hdqtrs., Saguaro Nat'l. Mon. East, 6-X-77 (CDJ #96-77); 31 mi NE Sells, 8-X-77 (CDJ #134-77).
- 3. Calliandra eriophylla Bentham: Arizona. Yavapai Co.: Black Canyon City, 30-V-76 (CDJ #2-76).
- 4. Lysiloma thornberi Britt. & Rose: Arizona. Pima Co.: ca 3700', Chimenea Cyn., Rincon Mts., 14-X-76 (CDJ #37-76).

Stator limbatus (Horn)

1. Acacia goldmanii (Br. & Rose) Wiggins: Mexico. Baja California Sur: 9 mi S Villa Constitucion, 25-XII-75 (CDJ #102-75).

2. Calliandra eriophylla Bentham: Arizona. Yavapai Co.: Black Canyon City,

30-V-76 (CDJ # #1-76 & 2-76).

3. Leucaena guatemalensis Britt. & Rose: Mexico. Colima: ca 1500', 3 mi S Co-

lima, 7-III-73 (CDJ #373-73).

4. Lysiloma thornberi Britt. & Rose: Arizona. Pima Co.: ca 3700', Chimenea Cyn., Rincon Mts., 14-X-76 (CDJ # #37-76 & 38-76).

Stator pruininus (Horn)

1. Acacia greggii Gray: Arizona. Yavapai Co.: Clear Creek Camp, 7 mi SE Camp Verde, 17-IX-77 (CDJ #87-77). Pima Co.: ca 2600′, 2.3 mi S Sells, 15-VIII-77 (CDJ #70-77).

2. Calliandra eriophylla Bentham: Arizona. Yavapai Co.: Black Canyon City,

30-V-76 (CDJ # #1-76 & 2-76).

Stator pygidialis (Schaeffer)

1. Calliandra reticulata Gray: Arizona. Cochise Co.: Naco, 7-X-77 (CDJ #116-77).

Stator sordidus (Horn)

1. Lysiloma divaricata (Jacq.) Macbride: Mexico. Sonora: 14 mi E Navojoa, 26-XII-76 (CDJ #191-76); 10 mi N Guaymas, 27-XII-76 (CDJ #203-76); 3 mi W Alamos, 27-XII-77 (CDJ #165-77); ca .5 mi E Alamos, 28-XII-77 (CDJ #176-77); 10 mi E Navojoa, 30-XII-77 (CDJ #201-77); 10 mi N Guaymas, 3-I-78 (CDJ #223-78).

2. Mimosa laxiflora Bentham: Mexico. Sonora: 10 mi E Navojoa, 30-XII-78 (CDJ

#236-78).

Stator vachelliae Bottimer

1. *Acacia cymbispina* Sprague & Riley: Mexico. Sonora: 13 mi NW Navojoa, 30-XII-77 & 26-XII-76 (CDJ # #208-77 & 197-76).

2. A. farnesiana (L.) Willd.: Mexico. Sonora: 11 mi S Hermosillo, 27-XII-76 (CDJ #212-76); 13 mi NW Navojoa, 30-XII-77 (CDJ #208-77).

Stator vittatithorax (Pic)

1. Lysiloma desmostachys Bentham: Mexico. Colima: ca 1100', 7 mi S Colima, 7-III-73 (CDJ #380-73).

Zabrotes interstitialis (Chevrolat)

1. Cassia moschata H.B.K.: Panama. Canal Zone: Ft. Clayton, 25-II-64 (LJB #121u).

Z. planifrons Horn

1. Cassia leptadenia Greenm: Arizona. Santa Cruz Co.: ca 4200', 6 mi E Ruby, 15-X-76 (CDJ #63-76).

Z. spectabilis Horn

1. Cassia roemeriana Scheele: Texas, Menard Co.: Menard, 13-14-VI-46 (LJB #86b).

Zabrotes sp. #1

1. Vicia leavenworthii Torr. & Gray: Texas: Bexar Co., 10-V-62 (LJB #112u).